

▶ **The Clinical Use of PET-CT in Radiotherapy Planning
Head and Neck Cancer**

V. Grégoire (Brussels)

The ultimate objective of radiotherapy is to achieve a high percentage of loco-regional control with a low incidence of morbidity, hence directly impacting on the overall survival and the quality of life.

In this framework, one of the first steps in the radiotherapy planning process is to precisely select and delineate the target volumes (i.e. the tumor) and the surrounding normal tissues potentially responsible for treatment morbidity if irradiated at a too high dose. For a long time, CT is used as the reference imaging modality as it can also be used for dose calculation (taking into account the Hounsfield units, which are surrogates of tissue density). This requires that the image acquisition is performed in treatment position, thus with the patients immobilized on a flat tabletop.

In pharyngo-laryngeal tumors, our group has shown that MRI did not bring any advantage over CT neither for the delineation accuracy, nor for the inter-observer variability. On the contrary, more recently, several groups including our, have shown that providing the images are acquired, reconstructed and segmented in a proper way, FDG-PET improved the delineation of pharyngo-laryngeal tumor volumes. This improvement translated into an improvement in dose distribution, i.e. a lower dose was delivered to the surrounding normal tissues. This finding opens the way for a possible increase in the dose prescription -thus potentially increasing the probability of loco-regional control- without increasing the dose to the surrounding normal tissues. Furthermore, the use of PET with other tracers imaging biological pathways involved in radiation response (e.g. cell proliferation, tumor hypoxia) open a new avenue to specifically deliver an extra dose to the PET-positive area, i.e. the so-called "dose painting" approach.

After a short description of the radiotherapy processes, the lecture will focus on the usefulness of the various anatomic and functional imaging modalities for radiotherapy treatment of head and neck cancer patients. Methodological aspects and pitfalls with the use of multimodality images will be highlighted.

References:

- Grégoire V, Langendijk JA, Nuyts S. Advances in Radiotherapy for Head and Neck Cancer. *J Clin Oncol.* 2015 Oct 10;33(29):3277-84.
- Leclerc M, Lartigau E, Lacornerie T, Daisne JF, Kramar A, Grégoire V. Primary tumor delineation based on (18)FDG PET for locally advanced head and neck cancer treated by chemo-radiotherapy. *Radiother Oncol.* 2015 Jul;116(1):87-93.
- Servagi-Vernat S, Differding S, Sterpin E, Hanin FX, Labar D, Bol A, Lee JA, Grégoire V. Hypoxia-guided adaptive radiation dose escalation in head and neck carcinoma: a planning study. *Acta Oncol.* 2015 Jul;54(7):1008-16.
- Grégoire V, Jeraj R, Lee JA, O'Sullivan B. Radiotherapy for head and neck tumours in 2012 and beyond: conformal, tailored, and adaptive? *Lancet Oncol.* 2012 Jul;13(7):e292-300.
- Daisne JF, Duprez T, Weynand B, Lonneux M, Hamoir M, Reychler H, Grégoire V. Tumor volume in pharyngolaryngeal squamous cell carcinoma: comparison at CT, MR imaging, and FDG PET and validation with surgical specimen. *Radiology.* 2004 Oct;233(1):93-100. Epub 2004 Aug 18. Erratum in: *Radiology.* 2005 Jun;235(3):1086.

Oct. 18